

Macroscopic Changes in the Human Placenta in Hypertensive Disorders of Pregnancy in Indian Women: A Cross-sectional Study

GAURAV KHASTGIR¹, VEENA VIDYA SHANKAR², RAJINI UDAY³, SHAILAJA SHETTY⁴

ABSTRACT

Introduction: Hypertension (HTN) is a common complication in 6-10% of pregnancies and is one of the prime factors for foetal, as well as, maternal fatalities and morbidities. Significant pathological changes are seen in the placenta in hypertensive pregnancies before affecting the foetus. These changes may serve as an early warning system for detecting foetal issues like low birth weight and growth restriction. Early detection and intervention of morphological changes in the placenta and umbilical cord can help prevent adverse foetal outcomes.

Aim: To study the macroscopic changes in the placenta in women with HTN induced by pregnancy compared to non hypertensive pregnant women.

Materials and Methods: The present cross-sectional descriptive study was conducted in the Department of Anatomy and the Department of Obstetrics and Gynaecology, Ramaiah Medical College Hospital, Bengaluru, Karnataka, India, between December 2017 and January 2019. A sample size of 43 each was taken for non hypertensive and hypertensive subjects, including those with preeclampsia and chronic HTN with superimposed preeclampsia. The placental morphological parameters, routine biochemical tests and neonatal outcomes of both groups were analysed statistically using the Shapiro-Wilk test, Independent

t-test and Chi-square test. A probability value less than 0.05 was considered for statistical significance.

Results: The mean age of participants was 28.35 ± 3.026 years in the hypertensive group and 26.23 ± 2.636 years in the control group. The circular, oval and irregular shapes were 29 (65.9%), 10 (22.7%) and 4 (9.3%) in non hypertensive subjects and 33 (78.6%), 7 (16.27%) and 3 (7.1%) in the hypertensive subjects, respectively. The surface area (mean- 142.44 ± 49.69 ; p-value <0.001), radius (mean- 6.622 ± 1.24 ; p-value <0.001), circumference (mean- 37.419 ± 6.091 ; p-value <0.001), the quantum of cotyledons (mean- 13.09 ± 3.344 ; p-value <0.001) were lesser in the hypertensive group than the non hypertensive subjects. There were differences in placental and foetal weight among non hypertensive and hypertensive subjects, but the differences were insignificant. The Appropriate for Gestational Age (AGA) babies were more in non hypertensive subjects.

Conclusion: The present study found hypertensive pregnancies to be associated with a greater occurrence of unfavourable foetal outcomes. Furthermore, it highlights the importance of investigating the placenta's physical attributes and their association with foetal outcomes to develop standard operating protocols for diagnostic and therapeutic approaches in normal/complicated pregnancies.

Keywords: Calcification, Eclampsia, Foetus, Pregnancy-induced hypertension, Preeclampsia

INTRODUCTION

Hypertension (HTN) is a common complication in 6-10% of pregnancies and is one of the prime factors for foetal, as well as, maternal fatalities and morbidities [1]. The placenta is a complex multifunctional organ of largely foetal origin with a significant role in foetal growth and development [2].

The health of the placenta is critical for a healthy baby. During pregnancy, the placenta facilitates haemopoiesis, foetal circulation, nutrition, removal of waste, hormone secretion, exchange of gases and immunity of the foetus [2]. The growth of the human foetus is limited by the inability of the placenta to supply nutrients and oxygen to the foetus, leading to foetal under-nutrition, poor growth and low birth weight [3]. It has been suggested that placental nutrient transfer is significantly affected by blood flow and morphological features like surface area and thickness of the placenta [4].

The placenta is currently the focus of interest because significant pathological changes are seen in it before affecting the foetus. Hence, the changes in the placenta may serve as "an early warning system" for foetal issues [5]. There are significant macroscopic placental changes resulting in foetal growth restriction and hypoxia [6]. Uteroplacental vascular thrombosis leads to placental ischaemia and accelerated placental maturation resulting in calcification. These are the most common gross morphological changes in severe Pregnancy-induced Hypertension (PIH) [7,8]. HTN is

associated with increased vascular resistance leading to decreased uteroplacental perfusion and foetal stem artery constriction resulting in an adverse foetal outcome [9]. All these changes result in a much smaller placental size. The placental weight and size are inversely proportional to the degree of PIH [7,9]. Studies have shown lower neonatal birth weight in PIH, and it has been positively correlated with placental weight. In addition, the mean placental surface area is also significantly lower with an inverse association between surface area and disease severity. Interestingly, the placental surface area also directly correlates with foetal distress [7,9]. In PIH, the umbilical cord diameter has been reported to be low. The site of cord insertion is more likely to be eccentric, central or marginal. This is thought to be associated with impaired placental perfusion leading to poor foetal outcomes and low birth weight [10-12]. By identifying morphological placental and umbilical cord changes, early intervention by delivery may prevent poor foetal outcomes.

Placental changes in women with hypertensive disorders have been studied. None of these studies have included the full spectrum of anatomical parameters or compared placental changes with all measures of foetal outcome. The primary aim was to study the macroscopic changes in the placenta in women with HTN induced by pregnancy compared to non hypertensive pregnant women. The objective was to associate the macroscopic placental changes with antenatal biochemical parameters and neonatal outcomes.

MATERIALS AND METHODS

A cross-sectional descriptive study was conducted in the Department of Anatomy and the Department of Obstetrics and Gynaecology, Ramaiah Medical College, Bengaluru, Karnataka, India, from December 2017 to January 2019. Prior ethical clearance was obtained from the Institutional Ethics Committee (IEC). Written informed consent was obtained from all participants.

Sample size calculation: Based on a previous study conducted by Udainia A and Jain ML, the mean placental weight was found to be 495 with a standard deviation of 114.11 in the control group as compared to 405.67 with a standard deviation of 101.64 in the PIH group [13]. In the present study, based on this finding, considering a similar mean difference and power of 90%, and an alpha error of 1%, the sample size was calculated to be 43 in each group.

Inclusion criteria: Inclusion criteria: Antenatal women presenting to the Labour Room of the study Institution both nulliparous and multiparous women, and with normal vaginal and caesarean section deliveries were included in the study.

Exclusion criteria: Women with maternal anaemia, gestational diabetes mellitus, multiple pregnancies, cardiac and renal diseases and cases with manual removal of retained placentae were excluded from the study.

Study Procedure

Placentae from the non hypertensive and hypertensive subjects were collected from the hospital, and the findings were compared with the patients' history and biochemical investigations during their routine antenatal visits. The haematological and biochemical reports were taken from the patients' files. Placentae were washed in tap water and blood clots were removed. The formaldehyde-preserved samples were examined for various parameters. The parameters observed were:

A. Maternal


1. **Blood pressure:** Based on the National High Blood Pressure Education Program's (NHBPEP) classification [14], the hypertensive subjects were grouped as given below:

- Group-1: Gestational HTN-BP $\geq 140/90$ mmHg for the first time during pregnancy; No proteinuria
- Group-2: Preeclampsia-BP $\geq 140/90$ mmHg after 20 weeks' gestation; proteinuria ≥ 300 mg/24 hours or $\geq 1+$ dipstick
- Group-3: Eclampsia-BP $\geq 160/110$ mmHg after 20 weeks' gestation; proteinuria ≥ 2000 mg/24 hours or $\geq 2+$ dipstick
- Group-4: Chronic HTN with superimposed preeclampsia- New onset proteinuria ≥ 300 mg/24 hours in hypertensive women but no proteinuria before 20 weeks' gestation
- Group-5: Chronic HTN-BP $\geq 140/90$ mmHg before pregnancy or diagnosed before 20 weeks' gestation or HTN first diagnosed after 20 weeks' gestation and persistent after 12 weeks postpartum

2. **Placental parameters:**

- Shape: circular, oval or irregular
- Attachment of umbilical cord: eccentric, marginal or central
- Weigh (in grams): measured using a digital weighing scale (before preserving the placentae in formalin solution).
- Diameter (in centimeters): Two diameters measured at right angles to each other. The mean of the two maximum diameters was considered. All measurements were performed by a single observer.
- The surface area (in cm^2) was calculated using $\pi(\text{radius})^2$

- Thickness (in centimeters): the maternal surface of the placenta was arbitrarily divided into three equal zones by drawing two circles. With a long needle, the thickness was measured at points 1-the center of the central zone; points 2a and 2b from the middle zone; points 3a and 3b-from the peripheral/outer zone. The mean of five measurements was considered for thickness [Table/Fig-1].



[Table/Fig-1]: Measurement of placental thickness.

- Mean quantum of cotyledons: on the maternal side of the placenta
- Presence or absence of calcification and retro placental clots.

3. **Biochemical parameters:**

- Serum Creatinine
- Platelet count
- Liver Function Test {Only Aspartate Aminotransferase (AST) and Alanine Transaminase (ALT)}. Parameters of the third trimester were taken for measurement.

B. **Foetal parameters:**

- Foetal weight-in grams
- Appearance, Pulse, Grimace, Activity and Respiration (APGAR) score at 1 and 5 minutes, respectively
- Foetal outcome
 - Appropriate for Gestational Age (AGA)
 - Stillbirth (SB)
 - Intrauterine Death (IUD)
 - Intrauterine Growth Retardation (IUGR)


STATISTICAL ANALYSIS

The data were analysed using the Statistical Package for Social Sciences (SPSS) software version 18.0. The normality of the data was analysed using the Shapiro-Wilk test. Descriptive statistics on macroscopic variables were described in terms of mean with standard deviation. The foetal outcome was summarised in terms of percentage. Independent t-test was used to analyse the differences in macroscopic changes in HTN and non hypertensive subjects. The Chi-square test was utilised to compare foetal outcomes in hypertensive and non hypertensive subjects. A probability value less than 0.05 was considered for statistical significance.

RESULTS

In the study, in Group-1, there were 24 samples; in Group-2, there were 13 samples; in Group-3, there were four samples; Group-4 had nil samples; and Group-5 had two samples. The mean age of participants was 28.35 ± 3.026 years in the hypertensive group and 26.23 ± 2.636 years in the control group.

In both hypertensive and non hypertensive subjects, the placenta was circular, oval and irregularly shaped. The circular, oval and irregular shapes were 29 (65.9%), 10 (22.7%) and 4 (9.3%) in non hypertensive subjects and 33 (78.6%), 7 (16.27%), and 3 (7.1%) in the hypertensive subjects, respectively [Table/Fig-2].



[Table/Fig-2]: Placental shapes and types of attachment of umbilical cord.


The current study showed a higher incidence of eccentric insertion, n=27 (61.4%) followed by central, n=12 (27.3%) and marginal, n=4 (9.3%) in the non hypertensive subjects. In comparison, insertion was central in 20 (46.5%), followed by marginal in 13 (30.1%) and eccentric in 10 (23.3%) hypertensive subjects. The p-value was <0.001 and hence, statistically significant.

Statistical significance was observed for placental radius, circumference, surface area and quantum of cotyledons. The difference in foetal weight was not significant between the two groups [Table/Fig-3].

Parameters	Non hypertensive subjects	Hypertensive subjects	p-value
Weight of placenta (grams)	515±142.16	484.88±151.99	0.342
Placental radius (cm)	7.613±0.96	6.622±1.24	<0.001*
Placental circumference (cm)	47.045±6.65	37.419±6.091	<0.001*
Placental surface area (cm ²)	184.90±45.40	142.44±49.69	<0.001*
Quantum of cotyledons in placenta	20.41±4.61	13.09±3.344	<0.001*
Umbilical cord thickness (cm)	5.98±1.79	5.21±1.46	0.32
Foetal weight (grams)	2581±63	2400±14	0.293

[Table/Fig-3]: Comparison of placental and foetal parameters. Fisher's-exact test and Mann-whitney test; *The p-value <0.05 was considered statistically significant

A higher incidence of calcified areas, 20 (46.5%) was noted in the hypertensive subjects; however, 2 (4.5%) were also seen in the non hypertensive subjects. Statistical significance was seen for calcified areas (p-value<0.001). Retroplacental clots were observed in 18 (41.86%) hypertensive and 10 (23.25%) non hypertensive subjects [Table/Fig-4].



[Table/Fig-4]: Retroplacental clots and calcified areas in the placenta. The red arrow is pointing towards the retroplacental clots and calcified areas, respectively

Among the biochemical parameters observed in the hypertensive group, platelet count, serum creatinine values, AST, and ALT levels were within the normal range in most of the hypertensive group [Table/Fig-5].

On comparing the foetal outcome, it was found that almost 38 (88.37%) of pregnancies in the non hypertensive subjects and 32 (74.41%) in the hypertensive subjects were AGA. The IUD was 1 (2.32%) and 2 (4.65%) in the non hypertensive and hypertensive subjects, respectively [Table/Fig-6].

It was observed that all 13 (100%) neonates with IUGR, 14 (20.28%) with AGA, and 3 (75%) with IUD/SB had a placental weight of less

Sl no	Biochemical parameters	Normal range	Hypertensive subjects, n (%)	Increased/Decreased
1	Platelet count (cells/mL)	1.5-4.5×10 ⁶ cells/L	41 (95.35)	Decreased in 2 (4.65%)
2	Serum creatinine (mg/dL)	0.5-0.09 mg/dL	41 (95.35)	Increased in 2 (4.65%)
3	AST (Units/Litre)	8-48 U/L	36 (83.73)	Increased in 7 (16.27%)
4	ALT (Units/Litre)	7-55 U/L	41 (95.35)	Increased in 2 (4.65%)

[Table/Fig-5]: Biochemical parameter values in hypertensive subjects.

Foetal outcome		Non hypertensive subjects, n (%)	Hypertensive subjects, n (%)	p-value
Appropriate for Gestational Age (AGA)	Yes	38 (88.37)	32 (74.41)	0.176
	No	5 (11.62)	11 (25.58)	
Stillbirth	Yes	0 (0)	1 (2.32)	0.314
	No	43 (100)	42 (97.68)	
IUD	Yes	1 (2.32)	2 (4.65)	0.557
	No	42 (97.68)	41 (95.35)	
IUGR	Yes	3 (6.98)	9 (20.93)	0.132
	No	40 (90.7)	34 (79.07)	

[Table/Fig-6]: Foetal outcome in the non hypertensive and hypertensive subjects.

than 450 grams, indicating poor foetal outcomes and supporting the well known theory of a small placenta being associated with a small baby. The p-value was not statistically significant (0.029) [Table/Fig-7].

Placental weight	Foetal outcome			Total N=86
	AGA n=69	IUGR n=13	IUD+SB n=4	
<450 gms	14 (20.28)	13 (100)	3 (75)	30 (34.88)
450-860 gms	55 (79.71)	0	1 (25)	56 (65.11)

[Table/Fig-7]: Placental weight and foetal outcome in both groups. Values are presented as n (%)

DISCUSSION

The placenta was circular, oval and irregular in both groups in the present study, similar to observations in studies reported by Goswami PR and Shah SN (2016), Kishwara S et al., (2009), Kumar VS et al., (2015), Shrestha I et al., (2022), and Wubale Y and Tolera A (2017), except in the study by Ranga MKSS et al., (2017) where all placentae in non hypertensive subjects were circular and in hypertensive subjects, the shape was circular followed by irregular with no ovoid-shaped placenta [9,15-19].

The site of attachment of the umbilical cord was central, eccentric and marginal in the non hypertensive subjects, as reported by Goswami PR and Shah SN (2016), Kaur P et al., (2014), Udainia A and Mehta CD, (2013), Abudullah AN et al., (2022), and Sarma GD et al., (2021), except in the present study and the one by Kumar VS et al., (2015) where the order of attachment was eccentric, central and marginal, respectively [9,12,16,20-22]. In the hypertensive subjects, the incidence of the site of cord insertion showed different patterns in various studies [9,12,16,20-23].

The placental thickness was less in the hypertensive group in various studies except the current study. The calcification was higher in the hypertensive subjects with respect to non hypertensive subjects in various studies [Table/Fig-8] [9,16,19,22-24].

The quantum of cotyledons in the present study was in line with those reported by Salmani D et al., (2014) in non hypertensive subjects, while in the hypertensive subjects, the cotyledons observed were more than the present study in other studies [Table/Fig-9] [11,15,16,18,19,25,26].

The placental surface area in the present study in the non hypertensive and hypertensive subjects was 184.90±45.40 cm² and 142.44±49.69 cm², respectively, which was low compared

Study	Placental thickness (in centimeters)		Placental calcification, n (%)	
	Non hypertensive subjects	Hypertensive subjects	Non hypertensive subjects	Hypertensive subjects
Kumar VS et al., (2015) [16]	3.17	2.28	28 (27.7)	34 (34)
Goswami PR and Shah SN (2016) [9]	2.60	2.20	8	18
Porwal V et al., (2017) [23]	Not a parameter		14 (46.66)	20 (66.66)
Ranga MKSS et al., (2017) [19]	2.4±0.1	1.9±0.9	7 (23.3)	17 (56.7)
Chhatwal J et al., (2018) [24]	2.10±0.52	1.91±0.47	12 (28.6)	19 (45.2)
Sarma GD et al., (2015) [22]	1.76±0.19	1.48±0.15	5	26
Present study	2.63±0.77	2.74 ±0.63	2 (4.5)	20 (46.5)

[Table/Fig-8]: Comparison of placental morphology in different studies [9,16,19,22-24].

Study	Quantum of cotyledons		p-value
	Non hypertensive subjects	Hypertensive subjects	
Majumdar S et al., (2005) [26]	17+2	16+2	>0.01
Kishwara S et al., (2009) [15]	15.77±2.80	14.30±2.47	<0.05
Salmani D et al., (2014) [25]	21.18±1.48	21.21±1.58	Not significant
Nahar L et al., (2015) [11]	17.40	15.39	<0.001
Kumar VS et al., (2015) [16]	18.54	17.12	<0.001
Ranga MKSS et al., (2017) [19]	19.1±1.2	16.1±2.5	<0.001
Wubale Y and Tolera A (2017) [18]	18.66±1.21	17.24±1.06	<0.001
Present study	20.41±4.61	13.09±3.344	<0.001

[Table/Fig-9]: Comparison of quantum of cotyledons in different studies [11,15,16,18,19,25,26].

to that observed by Sirpurkar M and Anjankar VP (2017) (223.6±108.5 and 188.4±89.6 in two groups, respectively) [27]. PIH significantly reduces placental weight and its dimensions and gives rise to characteristic histological changes. All these factors are responsible for ischaemia and the compromised uteroplacental blood flow leading to low foetal weight [28].

The placental circumference in the current study in the non hypertensive and hypertensive subjects was 47.045±6.65 cm and 37.419±6.091 cm, respectively, which was slightly less than that observed in the study by Kumar VS et al., (2015) in non hypertensive and hypertensive subjects with 45.1 cm and 41.1 cm, respectively [16].

The weight of the placenta and the foetal weight in the hypertensive subjects were less than in the non hypertensive subjects in the current and various studies [Table/Fig-10] [13,16,19,22,25-27,29-32].

The AGA in the non hypertensive subjects in the present study and that reported by Kaur P et al., (2014) and Sirpurkar M and Anjankar VP (2017) was in the range of 88.37% to 100% [12,27]. The IUGR (9.3%) and IUD (2.3%) in the non hypertensive subjects were

observed only in this study. Sirpurkar M and Anjankar VP (2017) have reported 3.3% SBs with none in the current study [27].

In the hypertensive subjects, the AGA observed was low (42.67%) in the study reported by Kaur P et al., (2014); ranged between 72.09% to 84.85% in the current study and that by Sirpurkar M and Anjankar VP (2017) [12,27]. A higher incidence of IUGR of 44% has been reported by Kaur P et al., (2014) and 20.93% in this study [12]. Sirpurkar M and Anjankar VP (2017) have reported 15.15% SBs, and in the current study, it is 2.3% [27]. In the present study, the differences between the non hypertensive and hypertensive subjects in parameters like placental radius, circumference, surface area and quantum of cotyledons were found to be statistically significant. The abnormal values of biochemical parameters and adverse foetal outcomes are higher in hypertensive subjects but need testing on a larger sample size to establish whether the differences are statistically significant.

Limitation(s)

A larger sample size in each group of hypertensive subjects needs to be tested for all parameters to establish statistically significant observations, which can lead to the categorisation of placental morphological changes in groups of hypertension (NHBPEP) classification.

CONCLUSION(S)

The hypertensive disorders of pregnancy adversely affect the morphological and morphometric features of the placenta, biochemical values such as serum creatinine, platelet count, AST and ALT levels, foetal weight, and foetal outcomes. The study calls attention to further research to establish the statistical significance of the above findings. It stresses the importance of placental examination and research by a trained professional for the proper identification of causes of poor pregnancy outcomes in hypertensive disorders of pregnancy to improve the modus operandi.

Study	Mean foetal weight (in grams)		Mean placental weight (in grams)	
	Non hypertensive subjects	Hypertensive subjects	Non hypertensive subjects	Hypertensive subjects
Udainia A and Jain ML (2001) [13]	2640.0	2280.0	495.0	405.67
Majumdar S et al., (2005) [26]	2800.0	2040.0	485.85	399.0
Salmani D et al., (2014) [25]	3140.0	2440.0	519.8	395.15
Singh S and Gugapriya TS (2014) [29]	3140.9	2329.72	435.92	345.96
Kumar VS et al., (2015) [16]	2556.93	2192.50	469.50	420.98
Londhe PS and Mane AB (2015) [30]	2730.0	2260.0	401.8	312.93
Gautam G et al., (2015) [31]	2634.59	2349.0	486.78	423.4
Sarma GD et al., (2015) [22]	2670.0	2400.0	514.8	486.9
Jain A et al., (2016) [32]	2633.6	2435.2	454.24	406.32
Ranga MKSS et al., (2017) [19]	2900	2500	516.7	455.7
Sirpurkar M and Anjankar VP (2017) [27]	2865	2267	486	398
Present study	2581	2400	515	484.88

[Table/Fig-10]: Comparison of foetal weight and placental weight in different studies [13,16,19,22,25-27,29-32].

Acknowledgement

The authors would like to acknowledge the Institute statisticians, Mr Shivaraj and Ms Radhika Haridas for their guidance.

REFERENCES

- [1] Kintiraki E, Papakatsika S, Kotronis G, Goulis DG, Kotsis V. Pregnancy-induced hypertension. *Hormones (Athens)* 2015;14(2):211-23.
- [2] Yetter JF. Examination of the placenta. *American Family Physician*. 1998;57(5):1045-54.
- [3] Kliman HJ. Behind every healthy baby is a healthy placenta. *Parenthood Lost*. Westport, CT: Bergin & Garvey. 2001;130-131.
- [4] Harding JE. The nutritional basis of the fetal origins of adult disease. *Int J Epidemiol*. 2001;30(1):15-23.
- [5] Fowden AL, Ward JW, Wooding FP, Forhead AJ, Constancia M. Programming placental nutrient transport capacity. *Physiol J*. 2006;572(1):05-15.
- [6] Benirschke K, Kaufmann P. *Pathology of the Human Placenta*. 2nd ed. New York: Springer Verlag; 1990.
- [7] Backes CH, Markham K, Moorehead P. Maternal preeclampsia and neonatal outcomes. *J Pregnancy*. 2011;2011:214365.
- [8] Udainia A, Bhagwat SS, Mehta CD. Relation between placental surface area, infarction and foetal distress in pregnancy induced hypertension with its clinical relevance. *J Anat Soc India*. 2004;53(1):27-30.
- [9] Goswami PR, Shah SN. Placenta in normal and pregnancy induced hypertension about its clinical significance: A gross study. *Int J Sci Stud*. 2016;4(7):58-61.
- [10] Garg K, Rath G, Sharma S. Association of birth weight, placental weight and the site of umbilical cord insertion in hypertensive mothers. *Journal of Anatomical Society of India*. 1996;44(4).
- [11] Nahar L, Nahar K, Hossain MI, Jahan S, Rahman MM. Placental changes in pregnancy-induced hypertension. *Mymensingh Medical Journal: MMJ*. 2013;22(4):684-93.
- [12] Kaur P, Kaushal S, Sharma S, Singh K. To study relationship of umbilical cord insertion with foetal outcome in pih and normotensive pregnancies. *Int J Pure App Biosci*. 2014;2(6):108-11.
- [13] Udainia A, Jain ML. Morphological study of placenta in pregnancy induced hypertension with its clinical relevance. *J Anat Soc India*. 2001;50(1):24-27.
- [14] Report of the National High Blood Pressure Education Program Working Group on high blood pressure in pregnancy. *Am J Obstet Gynecol*. 2000;183(1):S1-S22.
- [15] Kishwara S, Ara S, Rayhan KA, Begum M. Morphological changes of placenta in preeclampsia. *Bangladesh Journal of Anatomy*. 2009;7:49-54.
- [16] Kumar VS, Rani A, De'souza AS, Babu P. Effect of pregnancy induced hypertension on the morphometry of placenta and weight of new born. *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)*. 2015;14(7):87-95.
- [17] Shrestha I, Gautam P, Khajju S. Morphological variations in placentas among deliveries in the department of obstetrics and gynaecology in a tertiary care centre: A descriptive cross-sectional study. *J Nepal Med Assoc*. 2022;60(253):808-10.
- [18] Wubale Y, Tolera A. Gross morphological study of placenta in preeclampsia mothers. *Anatomy Journal of Africa*. 2017;6(2):977-81.
- [19] Ranga MKSS, Kumar KV, Thangam A, Mallika MCV. Morphological variations of human placentae in preterm labor, pregnancy-induced hypertension and gestational diabetes mellitus. *Int J Sci Stud*. 2017;4(11):144-46.
- [20] Udainia A, Mehta CD. Study of umbilical cord in pregnancy induced hypertension. *National Journal of Medical Research*. 2013;3(1):66-69.
- [21] Abudullah AN, Al-Bakri NA, Abdulhamed WA. Histopathological changes in placentas of gestational hypertension women. *Int. J. Drug Deliv. Technol*. 2022;12(1):256-59.
- [22] Sarma GD, Sanyal R, Al Hossaini SS, Khemka VK, Mitra A. Study the morphology and the morphometric measurements of placenta among normotensive and pregnancy-induced hypertension pregnancies. *International Journal of Health and Clinical Research*. 2021;4(7):142-47.
- [23] Porwal V, Jain D, Gupta S, Khandelwal S, Kasliwal N. Spectrum of placental changes in pregnancy induced hypertension. *Annals of Pathology and Laboratory Medicine*. 2017;4(1):A70-A76.
- [24] Chhatwal J, Chaudhary DN, Chauhan N. Placental changes in hypertensive pregnancy: A comparison with normotensive pregnancy. *Int J Reprod Contracept Obstet Gynecol*. 2018;7(9):3808-13.
- [25] Salmani D, Purushothaman S, Somashekara SC, Gnanagurudasan E, Sumangaladevi K, Harikishan R, et al. Study of structural changes in placenta in pregnancy-induced hypertension. *J Nat Sc Biol Med*. 2014;5(2):352-55.
- [26] Majumdar S, Dasgupta H, Bhattacharya K, Bhattacharya A. A study of placenta in normal and hypertensive pregnancies. *J Anat Soc India*. 2005;54(2):01-09.
- [27] Sirpurkar M, Anjanekar VP. Study of correlation between placental morphology and adverse perinatal outcome in different conditions affecting pregnancy. *Int J Reprod Contracept Obstet Gynecol*. 2015;4(4):1165-68.
- [28] Mandhana VS, Shroff GA, Kharkar AR, Nawal A. A comparative morphological study of placenta in normal pregnancy and placenta of pregnancy induced hypertension. *International Journal of Health Sciences and Research (IJHSR)*. 2014;4(12):184-88.
- [29] Singh S, Gugapriya TS. A cross-sectional morphometric study of hypertensive with normal placentae and its correlation with fetal outcome. *Int J Anat Res*. 2014;2(2):437-42.
- [30] Londhe PS, Mane AB. Morphometric study of placenta and its correlation in normal and hypertensive pregnancies. *International Journal of Pharma and Bio Sciences*. 2011;2(4):429-37.
- [31] Gautam G, Bhake A, Samal N, Shukla S, Gautam M, Patil P. Correlative study of placenta in Pregnancy induced hypertension cases. *International Journal of Development Research*. 2015;5(6):4802-06.
- [32] Jain A, Sharma D, Sharma V, Baweja S, Jain R. Study of placental weight in normal and pre-eclamptic pregnancies and its correlation with birth weight. *Int J Med Res Rev*. 2016;4(4):562-68.

PARTICULARS OF CONTRIBUTORS:

1. Junior Resident, Department of Obstetrics and Gynaecology, PGIMER, Chandigarh, India.
2. Professor, Department of Anatomy, Ramaiah Medical College, Bengaluru, Karnataka, India.
3. Professor, Department of Obstetrics and Gynaecology, Ramaiah Medical College, Bengaluru, Karnataka, India.
4. Professor, Department of Anatomy, Ramaiah Medical College, Bengaluru, Karnataka, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Gaurav Khastgir,
Junior Resident, Department of Obstetrics and Gynaecology, Room Number-10,
PGIMER Chandigarh Sector 12, Chandigarh-160012, India.
E-mail: gauravkhastgir1996@gmail.com

PLAGIARISM CHECKING METHODS: [\[Jain H et al.\]](#)

- Plagiarism X-checker: Feb 12, 2024
- Manual Googling: Apr 24, 2024
- iThenticate Software: Apr 29, 2024 (7%)

ETYMOLOGY: Author Origin

EMENDATIONS: 6

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. Yes

Date of Submission: **Feb 12, 2024**

Date of Peer Review: **Mar 20, 2024**

Date of Acceptance: **Apr 30, 2024**

Date of Publishing: **Jul 01, 2024**